

Masses of Quarks and Leptons in a Supersymmetric GUT.

B. B. Deo, L. Maharana^{1,*} and P. K. Mishra²

¹*Department of Physics, Utkal University, Bhubaneswar-751004, India.*

²*Department of Physics, S.C.S. College, Puri-752001, India*

(Dated: July 18, 2009)

The masses and their variation with energy of the elementary particles obtained by solving the one loop renormalization group equation (RGE) in the Minimal Supersymmetric Standard Model (MSSM) is considered. A mass about 115 GeV for all the fermions at GUT scale is found as the possible solution of RGE. The mass spectra of all the fermions are also obtained within acceptable errors. They are shown graphically to descend for the top quark but ascend for all other fermions from the laboratory to the GUT scale.

PACS numbers: 12.10.Dm, 12.10.Kt

Keywords: RGE, MSSM

Only one of all the parameters of the standard model was first predicted in 1974 by Gaillard and Lee[1]. Thereafter came the work of the mass matrix ansatz of Fritzsch [2]. A complete listing of textures and their relevance to experimental findings were made by Ramond, Roberts and Ross (RRR)[3]. Regarding the mass of the top quark, Pendleton and Ross[4], Faraggi[5] and some others [6, 7, 8, 9] have predicted the value to be around 175 GeV, even before the top was discovered. Here, however, we choose $M_{top}(M_X) = M_F(X) = M_U \simeq 115$ GeV as the only input.

The gauge sector of the standard model is characterised by three coupling constants g_3, g_2 and g_1 of $SU_C(3) \otimes SU_L(2) \otimes U_Y(1)$, respectively. For the three couplings at mass $M_X = 2.2 \times 10^{16}$ GeV[10] unite to a unified coupling constant $\frac{g_U^2}{4\pi} = \frac{1}{24.6}$. The RG equations for the couplings in the lowest order are given by

$$16\pi^2 \frac{dg_i(t)}{dt} = c_i g_i^3(t), \quad i = 1, 2, 3. \quad (1)$$

The coefficients are $c_1 = 6.6, c_2 = 1, c_3 = -3$.

The running parameter t is defined as $t = \log_e \frac{\mu}{M_Z}$, so that it varies from 0 to $\log_e \frac{M_X}{M_Z} \simeq 33$. The solution of equation(1) is

$$\frac{4\pi}{g_i^2(t)} = \frac{4\pi}{g_i^2} - \frac{c_i}{2\pi} t \quad (2)$$

Here, $g_i^2 = g_i^2(0)$ are the coupling strengths in the electroweak scale M_Z . Taking the value of $M_X = 2.2 \times 10^{16}$ GeV and $\frac{4\pi}{g_U^2} = 24.6$, we calculate the value of $\frac{4\pi}{g_1^2} = 59.24$, $\frac{4\pi}{g_2^2} = 29.85$ and $\frac{4\pi}{g_3^2} = 8.85$. These are consistent with experimental results. Thus the three gauge couplings are descendants of one coupling constant g_U .

We write the Yukawa sector SUSY RG equations, which had also been written by Babu[11] following Georgi and Glashow, and Cheng et al[12], with the masses in units of 175 GeV, which is taken to be the mass of the top quark,

$$16\pi^2 \frac{dM_F(t)}{dt} = A_F M_F^3(t) + [Y_F(t) - G_F(t)] M_F(t) \quad (3)$$

$$= A_F M_F^3(t) + Z_F(t) M_F(t), \quad (4)$$

where A_F is 6 for quarks i.e. $F=1, \dots, 6$ and it is 4 for leptons i.e. $F=7, 8, \dots, 12$. The way, the RGE written above, is critical and crucial. The first term is like the first term of gauge sector and $Z_F = Y_F(t) - G_F(t)$. Here the mass in M_F of equation (4) missing and this mass in M_F behaves as an external field like the problems of quantum mechanics Y_F is a mixed matrix term,

$$Y_F(t) = \sum_H A_{FH} M_H^\dagger(t) M_H(t), \quad H = 1, 2, 3, \dots, 12. \quad (5)$$

*Electronic address: lmaharan@iopb.res.in

In MSSM, this matrix A_{FH} is given by the 144 elements given below,

$$A_{FH} = \begin{pmatrix} 0 & 3 & 3 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 3 & 0 & 3 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 3 & 3 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 0 & 0 & 0 & 3 & 3 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 3 & 0 & 3 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 3 & 3 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 3 & 3 & 3 & 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 3 & 3 & 3 & 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 3 & 3 & 3 & 1 & 1 & 0 & 0 & 0 & 1 \\ 3 & 3 & 3 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 1 \\ 3 & 3 & 3 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 0 & 1 \\ 3 & 3 & 3 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 \end{pmatrix}. \quad (6)$$

The gauge factor $G_F(t)$, which are the sum of gauge couplings, are fixed. We take the values from references [13] and [14].

$$G_U(t) = \frac{13}{15}g_1^2(t) + 3g_2^2(t) + \frac{16}{3}g_3^2(t) = \sum_{i=1}^3 K_U^i g_i^2(t). \quad (7)$$

$F = 1, 2, 3$ stand for U and they are degenerate electromagnetic gaugewise. Similarly,

$$G_D(t) = \frac{7}{15}g_1^2(t) + 3g_2^2(t) + \frac{16}{3}g_3^2(t), \quad F = 4, 5, 6, \quad (8)$$

$$G_E(t) = \frac{9}{5}g_1^2(t) + 3g_2^2(t), \quad F = 7, 8, 9, \quad (9)$$

and

$$G_N(t) = \frac{3}{5}g_1^2(t) + 3g_2^2(t), \quad F = 10, 11, 12. \quad (10)$$

Here, $K_N^3 = K_E^3 = 0$, as the leptons do not have the strong colour interaction. We shall need the integrals

$$-\frac{1}{8\pi^2} \int_0^t d\tau G_F(\tau) = \sum_{i=1}^3 \frac{K_i^F}{c_i} \log \left(1 - \frac{c_i g_i t}{8\pi^2} \right), \quad (11)$$

and

$$-\frac{1}{8\pi^2} \int_0^{t_X} d\tau G_F(\tau) = \sum_{i=1}^3 \frac{K_i^F}{c_i} \log \left(1 - \frac{c_i g_i t_X}{8\pi^2} \right) = \sum_{i=1}^3 \frac{K_i^F}{c_i} \log \frac{g_i^2}{g_U^2}. \quad (12)$$

Deo and Maharana[15], and Deo, Maharana and Mishra[16] observed that equation (4), which has to be solved for a given fermion, does not contain the coefficients of the same mass in the matrix A . This is the reason we can integrate the RGE. This fact has probably been overlooked by previous authors, so their result, with the calculational details become cumbersome and the values obtained take enormous amount of computation.

Thus the present approach gives a simple method of entangling the mass due to finding the solution of 12 differential equations. The terms $Z_F(t)$ can be exponentiated away. We introduce a subsidiary mass $m_F(t)$ through

$$M_F(t) = m_F(t) \exp \left(\frac{1}{16\pi^2} \int_0^t Z_F(\tau) d\tau \right), \quad (13)$$

such that $M_F(M_Z) = m_F(M_Z) = m_F(0)$. They satisfy the equation

$$16\pi^2 \frac{dm_F(t)}{m_F^3} = A_F \exp \left(\frac{1}{8\pi^2} \int_0^t Z_F(\tau) d\tau \right) dt. \quad (14)$$

They look similar to the gauge sector RG equation (1). This can be solved exactly and integrating equation(14) from M_Z to M_X , i.e., from $t = 0$ to $t = t_X$, we get

$$\frac{8\pi^2}{m_F^3(M_Z)} = \frac{8\pi^2}{m_F^3(M_X)} + A_F \int_0^{t_X} dt \exp \left(\frac{1}{8\pi^2} \int_0^t Z_F(\tau) d\tau \right). \quad (15)$$

Putting back the exponential, we get

$$\frac{M_{top}^2(M_Z)}{M_F^2(M_Z)} = \frac{M_{top}^2(M_Z)}{M_F^2(M_X)} \exp \left(\frac{1}{8\pi^2} \int_0^{t_X} Z_F(\tau) d\tau \right) + \frac{A_F}{8\pi^2} \int_0^{t_X} dt \exp \left(\frac{1}{8\pi^2} \int_0^t Z_F(\tau) d\tau \right). \quad (16)$$

This is the exact one loop solution in this approximation. $M_F^2(M_Z)$ is the square of the mass of the fermions at M_Z . The descent or ascent running of the masses from GUT M_X to electroweak M_Z , can be obtained by integrating equation (14) from $t = t_X$ to t . The result which is not given in reference[15] is

$$\frac{8\pi^2 M_{top}^2}{M_F^2(t)} = \frac{8\pi^2 M_{top}^2}{M_F^2(M_X)} \exp \left(\frac{1}{8\pi^2} \int_t^{t_X} Z_F(\tau) d\tau \right) + A_F \int_t^{t_X} dt_1 \exp \left(\frac{1}{8\pi^2} \int_t^{t_1} Z_F(\tau) d\tau \right). \quad (17)$$

This is also one loop exact in this approach. By solving equations(16) and (17), we can find $M_F(M_X)$ and $M_F(t)$ respectively.

One retains only those terms containing $M_{top}^2(t)$ occuring in any integral with $Y_F(t)$ in the first approximation. For the top case, $Y_{top}(t)$ can be set equal to zero in the RG equation for the top is a very good approximation. The top mass is then given by,

$$1 = \frac{M_{top}^2(M_Z)}{M_{top}^2(M_X)} C_{top} + D_{top}, \quad (18)$$

where, using integrals(11) and (12)

$$C_{top} = \prod_{i=1}^3 \left(\frac{g_i^2}{g_U^2} \right)^{\frac{\kappa_i^F}{c_i}} = 0.086, \quad (19)$$

and

$$D_{top} = \frac{6}{8\pi^2} \int_0^{t_X=33} dt \prod_i^3 \left(1 - \frac{g_i^2 c_i t}{8\pi^2} \right)^{\frac{\kappa_i^F}{c_i}} = 0.802. \quad (20)$$

Putting these values, the original mass of the top, at an energy of 2.2×10^{16} GeV is,

$$M_{top}(M_X) = M_{top}(M_Z) (1 - D_{top})^{-\frac{1}{2}} C_{top}^{\frac{1}{2}} = 114 \text{ GeV}. \quad (21)$$

This has been Deo-Maharana's [15] result.

The top mass decreases with energy but all the other quarks and leptons, starting from the value at M_X acquire smaller values and become quite light in the electroweak scale. The descent or ascent equation(17) describing the 'run' can be put in a form like equation

$$\frac{M_{top}^2(M_Z)}{M_F^2(t)} = \frac{M_{top}^2(M_Z)}{M_F^2(M_X)} C_F(t) + D_F(t), \quad (22)$$

where

$$C_F(t) = a_F(t) \exp \left(\frac{1}{8\pi^2} \int_0^{t_X} Y_F(\tau) d\tau \right)$$

and

$$D_F(t) = \frac{A_F}{8\pi^2} \int_0^{t_X} dt_1 b_F(t_1) \exp \left(\frac{1}{8\pi^2} \int_t^{t_1} Y_F(\tau) d\tau \right).$$

Here,

$$a_F(t) = \prod_{i=1}^3 \left[\frac{\left(1 - \frac{g_i^2 c_i t_X}{8\pi^2}\right)}{\left(1 - \frac{g_i^2 c_i t}{8\pi^2}\right)} \right]^{\frac{K_i^F}{c_i}}, \quad (23)$$

and

$$b_F(t) = \prod_{i=1}^3 \left[\frac{\left(1 - \frac{g_i^2 c_i t_1}{8\pi^2}\right)}{\left(1 - \frac{g_i^2 c_i t}{8\pi^2}\right)} \right]^{\frac{K_i^F}{c_i}}. \quad (24)$$

This is the general result for all the masses.

For the top, we can take $Y_F \rightarrow 0$ and calculate the variation of its mass from $M_U \simeq 115$ GeV to the top mass 175 GeV using the equation (25) below

$$M_{top}(t) = \frac{M_{top}(M_X)}{\left[C_{top}(t) + \frac{M_U^2}{M_{top}^2} D_{top}(t) \right]^{1/2}}. \quad (25)$$

The matrix A_{FH} of equation(6) gives

$$Y_{top}(t) = 3M_c^2(t) + 3M_u^2(t) + M_b^2(t) + M_{\nu_\mu}^2(t) + M_{\nu_e}^2(t) + M_{\nu_\tau}^2(t). \quad (26)$$

The equation for the top becomes

$$16\pi^2 \frac{dM_{top}(t)}{dt} = 6M_{top}^3 + [Y_{top}(t) - G_{top}(t)] M_{top}(t). \quad (27)$$

For bottom, however,

$$16\pi^2 \frac{dM_{bottom}(t)}{dt} = 6M_{bottom}^3 + [Y_{bottom}(t) - G_{bottom}(t)] M_{bottom}(t). \quad (28)$$

M_{bottom} is considerably smaller than M_{top} . Furthermore,

$$Y_{bottom}(t) = M_{top}^2(t) + 3M_c^2(t) + 3M_s^2(t) + M_\mu^2(t) + M_e^2(t) + M_\tau^2(t). \quad (29)$$

This type of ordering approaches fails to solve the problem of all other fermions. So we find a different method presented below to obtain the values of masses and their variation with energy. Let us construct another function for fermions other than the top $B_F(t)$ and $B_{top}(t)$ such that

$$8\pi^2 \frac{d}{dt} \log B_F^2(t) = Z_F(t). \quad (30)$$

In particular,

$$\exp\left(\frac{1}{8\pi^2} \int_0^{t_X} Z_F(\tau) d\tau\right) = \frac{B_F^2(M_X)}{B_F^2(M_Z)} \quad \text{and} \quad \exp\left(\frac{1}{8\pi^2} \int_0^t Z_F(\tau) d\tau\right) = \frac{B_F^2(t)}{B_F^2(M_Z)}. \quad (31)$$

Using equations (16) and (30), we find,

$$M_F(M_Z) \simeq M_F(M_X) \frac{B_F(M_Z)}{B_F(M_X)} = M_F(M_X) \exp\left(\frac{-I_F}{16\pi^2}\right). \quad (32)$$

The integral I_F is

$$\begin{aligned}
I_F &= \int_0^{t_X} Z_F(t) dt = \int_0^{t_X} \left(\sum_G A_{FG} M_G^\dagger(t) M_G(t) - G_F(t) \right) dt, \\
&= \frac{1}{2} \int_0^{t_X} \left(\sum_G A_{FG} M_G^\dagger(t) M_G(t) - G_F(t) + \sum_G A_{FG} M_G^\dagger(-t) M_G(-t) - G_F(-t) \right) dt, \\
&= \frac{1}{4} \int_{-t_X}^{t_X} \frac{dt}{dM_F} dM_F \left(\sum_G A_{FG} M_G^\dagger(t) M_G(t) - G_F(t) + \sum_G A_{FG} M_G^\dagger(-t) M_G(-t) - G_F(-t) \right), \\
&= \frac{16\pi^2}{4} \int_{-M_U}^{M_U} \frac{dM_F(t)}{M_F} \frac{\left(\sum_G A_{FG} M_G^\dagger(t) M_G(t) - \frac{1}{2} (G_F(t) + G_F(-t)) \right)}{\left(A_F M_F^\dagger(t) M_F(t) - G_F(t) \right)}. \tag{33}
\end{aligned}$$

Here we have used equation(4). M_U is defined below. The poles in the integrand as can be easily noticed. Set

$$M_F(t) = M_U e^{i\theta_F(M_F(t))n_F} = M_U e^{i\theta_F(t)n_F},$$

where n_F and θ_F specify a particular Riemann sheet (This M_U should not be confused with the mass of the U-quark). n_F is a rotational integer. $t = M_U e^{i\theta_F(t)n_F}$ maps the complex t -plane to the inside of a circle of radius M_U . Using this in equation(33), we get

$$I_F = i \frac{16\pi^2}{2} \oint n_F d\theta_F(t) \frac{\sum_G A_{FG} - \frac{G_F(t)}{M_U^2}}{A_F + \sum_G A_{FG} - \frac{G_F(t)}{M_U^2}}. \tag{34}$$

There are poles in the integrand depending on M_U and G_F . The factor G_F/M_U^2 in the numerator is inconsequential and need be dropped. For the quarks, $A_F + \sum_{G=1}^6 = 6 + 7 = 13$, for the leptons $4 + 9 = 13$ and the integral is nearly a constant. The integer n_F , characterising F from the integral, using $1 = \frac{1}{12} \sum_H$ in equation(34) is in order and we find

$$\bar{I}_F = i n_F \frac{16\pi^2}{2} \frac{1}{12} \sum_H \sum_G A_{AG} \int_{-t_X}^{t_X} d\theta_H(t) \frac{1}{A_H + \sum_G A_{HG} - \frac{G_H}{M_U^2}}. \tag{35}$$

In the above, one need to take the averaged $\bar{I}_F$ value over the twelve fermions. Retracing the steps and using $M_H(t) = M_U e^{i\theta_H(t)n_F}$ in equation(34), we finally get

$$\bar{I}_F \simeq n_F \frac{1}{12} \sum_H \sum_G A_{HG} \int_0^{t_X} dt = \frac{1}{12} n_F t_X \sum_H \sum_G A_{HG}. \tag{36}$$

As G vary from 1 to 6, whereas F will vary from 2 to 12. Only the coefficients A_{HG} are needed to calculate $\bar{I}_F$ of equation(36). The coefficients A_{FG} have nonvanishing values for $F = 2, 3, \dots, 12$ and $G = 1, 2, \dots, 6$.

$$\text{For } F = 2, 3, \dots, 6: \quad A_{F_1} + A_{F_2} + \dots + A_{F_6} = 7, \quad \text{and for } F = 7, 8, \dots, 12: \quad A_{F_1} + A_{F_2} + \dots + A_{F_6} = 9, \tag{37}$$

For the twelve fermions, the sum of the coefficients A_{FG} is $(7 \times 5) + (9 \times 6) = 89$. The average of Z , i.e., $\bar{Z}$ is given by

$$Z = \frac{89}{12} \quad \text{and} \quad \bar{I}_F = n_F t_X \frac{89}{12}. \tag{38}$$

The masses of all the fermions, due to quark-lepton equivalence, other than the top is

$$M_F(M_Z) \simeq M_F(M_X) e^{-n_F \frac{t_X}{16\pi^2} \frac{89}{12}} \simeq \lambda^{n_F} M_U, \tag{39}$$

where $\lambda = \exp\left(-\frac{t_X}{16\pi^2} \frac{89}{12}\right) = 0.219$. This is the Wolfenstein parameter and is an excellent result in spite of the approximate estimates. The values of n_F as in the Table-I for leptons are 8 for electron(e), 4 for muon(μ), 3 for tau(τ), 16 for electron neutrino(ν_e), 9 for muon neutrino(ν_μ), 6 for tau neutrino(ν_τ) and that of quarks are 7 for up(u), 6 for down(d), 4 for strange(s), 3 for charm(c) and 2 for bottom(b).

TABLE I: Identification of Fermions(Leptons and Quarks)

n_F	Mass(GeV)	Fermion(Quarks and Leptons)	Expt.(GeV)
2	5.5	bottom(b)	
3	1.2	charm(c),tau lepton(τ)	1.4, 1.7
4	0.264	strange(s), muon(μ)	0.1 - 0.23, 0.01
6	0.012	down(d),tau neutrino(ν_τ)	0.055 - 0.115
7	0.0028	up(u)	0.003
8	6×10^{-4}	electron(e)	$\times 10^{-4}$
9	1.33×10^{-4}	muon neutrino(ν_μ)	1.9×10^{-4}
16	3×10^{-9}	electron neutrino(ν_e)	3×10^{-9}

It is important to note that because of equation(16), the variation of mass with energy comes out automatically also. With $Z_F(t) = 8\pi^2 \frac{d}{dt} \log B_F^2(t)$, this reduces to

$$\frac{M_{top}^2}{M_F^2(t)} = \frac{M_{top}^2}{M_U^2} \frac{B_F^2(t_X)}{B_F^2(t)} + \frac{A_F}{8\pi^2} \int_t^{t_X} \frac{B_F^2(t_1)}{B_F^2(t)} dt_1. \quad (40)$$

An approximate form of this is to neglect the very small contribution from the A_F and we get

$$M_F(t) \simeq M_U \lambda^{n_F(1-t/33)}. \quad (41)$$

This is shown in the Figure-1 which shows the variations explicitly. Much more exact analytic studies are needed to deduce the exact mass values from the RGE. This method does not depend on any previous calculation which do not take both quarks and leptons. We have succeeded in finding the correct values of the masses and the running masses of the entire Yukawa masses and we believe that these results will form an important role in the physics of particles.

We are very much thankful Dr P K Jena for his help.




FIG. 1: Variation of Masses(GeV) of all 12 fermions with t/t_X , $t=\log(\mu/M_Z)$ and $t_X=33$.

-
- [1] M.K.Gaillard, B.W.Lee, Phys. Rev. **D10**, 897 (1974)
 - [2] H.Fritzsch, Phys.Lett. **B70**, 437 (1977)
 - [3] P.Ramond, R.G.Roberts, G.G.Ross, Nucl. Phys. **B406**, 19 (1993)
 - [4] B. Pendleton, G. G. Ross, Phys. Lett. **B98**, 291 (1981)
 - [5] A. E. Faraggi, Phys. Lett. **B274**, 47 (1992)
 - [6] C. R. Das, M. K. Parida, hep-ph/0010004;
 - [7] M K Parida, B Purukayasth, hep-ph/9902374
 - [8] D.A.Demir, Renormalisation Group Invariants and Sparticle spectroscopy in the MSSM, e-print hep-ph/0408043
 - [9] B.Grzadkowski, M.Lindner, S.Theisen, Phys. Lett. **B198**, 64 (1987),
M.Olechowski, S.Poroski, Phys. Lett. **B257**, 388 (1991)
 - [10] M. B. Einhorn, D. R. T. Jones, Nucl. Phys. **B196**, 475 (1982),
C. G. Ross, R. G. Roberts, Nucl. Phys. **B377**, 574 (1992)
 - [11] K. S. Babu, Z. Phys., **C35**, 69 (1987)
 - [12] H. Georgi, S. L. Glashow, Phys. Rev. Lett. **32**, 438 (1974),
T. P. Cheng, E. Eichten, L. F. Lie, Phys. Rev. **D9**, 2259 (1974)
 - [13] R. Gotto, G. Sartori, M. Tonin, Phys. Lett. **B28**, 128 (1968),
R. J. Oakes, Phys. Lett. **B29**, 683 (1969); **B30**, 262 (1970)
 - [14] S. Dimopoulos, L.J.Hall, S.Raby, Phys. Rev. **D45**, 4192 (1992)
 - [15] B.B.Deo, L.Maharana, Phys. Lett. **B597**, 192 (2004),
 - [16] B.B.Deo, L.Maharana, P.K.Mishra, Phys. Lett. **B632**, 695 (2006)